Site: A.L. Tay Ist Break: 2.8 Other:

VALLEY OF THE DRUMS

Shepherdsville, Kentucky

MAFI

Spring 1979



DRAFT

Prepared by:

Royal J. Nadeau
U.S. Environmental Protection Agency
Environmental Response Team
Edison, New Jersey









# ACKNOWLEDGEMENT

This report would not be possible without the cooperative efforts of numerous individuals associated with Region IV's Emergency Response Branch, Water Surveillance Branch, Technical Support Branch and Ecology Branch plus the Environmental Response Team and the Kentucky Division of Water Resources.

#### A.L. TAYLOR DUMP SITE (VALLEY OF THE DRUMS)

# INTRODUCTION

On March 2, 1979, Region IV activated the Revolving Fund to clean up a spill from the Valley of the Drums in Bullit County, Kentucky. Jack Stonebreaker assumed the role of OSC and directed the contractor, OH Materials, EPA's Environmental Response Team, Coast Guard's Gulf and Pacific Strike Team, the Coast Guard's PIA Team, U.S. Fish and Wildlife and the State of Kentucky in a coordinated effort to contain and mitigate the environmental emergency at the Valley of the Drums.

The dump was opened around 1964 and operated by Mr. A.L. Taylor. In 1977, the dump was closed and Mr. Taylor died in early 1978. During the operation of the facility, Mr. Taylor never obtained a permit.

Records obtained from industrial firms contrast from records obtained from the Taylor Company. Taylor's records are incomplete especially regarding a period of heavy activity during 1975 and 1976.

Industrial records indicate a total of 27,389 drums were delivered to the site while Taylor's records show only 7,339 being delivered to the site. Industries that have provided the state with records are: Reliance Universal, Inc. (13,301); Ford Motor Company (10,105); George W. Whitesides Company (3,082); Kurfees Coating (489); Tremco (200); C&C Supply (98); Guardsman Chemical (67); Louisville Varnish (33); and Randal VeVay (14).

Assuming the total of 27,389 drums and the above ground count of 16,383 drums, the remaining 11,006 drums would approximate the number of buried drums on site. This number, however, is based on records from selected industries and would not indicate drums from other sources.

#### ENVIRONMENTAL INVESTIGATION

A series of field sampling efforts were conducted by EPA and State of Kentucky Water Resources personnel to document the occurrence and extent of contamination in the surface water and soils on the site proper and adjacent receiving waters. The initial sampling was performed to determine oil and grease, the presence of which in the adjacent waters was justification for the use of 311(k) funds for mitigation and cleanup.

#### EPA/REGION IV AND ERT SAMPLING EFFORTS

The first environmental samples in connection with the A.L. Taylor

Site (Valley of the Drums) was collected by Region IV Surveillance and

Analysis personnel on February 13, 1979 ( ). On this date, water

and sediment samples were collected from vilson Creek at sites downstream from

Valley. This was part of a larger sampling effort that was being performed in

connection with other hazardous waste disposal sites in and around the

Louisville area that were being investigated.

In the latter part of February (February 22), site specific sampling was performed on the A.L. Taylor site to delineate the types and quantities (see Figure 1 in Appendix).

of hazardous materials presents At this time, water samples were collected from standing water pools and melt water running off the properties of Wilson Creek.

Also, sediment samples were collected at specific sites to determine the presence of contamination in the surface soils.

In early March, during the early stages of cleanup, water and soil samples were collected by John Gilbert, EPA-ERT and Environmental Consultants, Inc., under contract to EPA. These samples were collected from specific areas within the A.L. Taylor properties and downstream in Wilson Creek.

On March 7th, an in-situ examination of Wilson Creek, the remote stream immediately adjacent to the Valley, were conducted by Don Schultz, Region IV Biologist and Royal J. Nadeau, ERT Biologist at the request of the OSC to perform an assessment of the biological and ecological state of the

stream (see attachment map in Appendix).

This assessment was performed by sampling and examining the benthic infauna of Wilson Creek. A small dip net (approximately 20 mm in diameter flattened on one side) was used to sample the infaunal community at various locations in the upper reaches of Wilson Creek. Most of the stations were downstream from the Valley of the Drums, with one being on a tributary (Station 2 Control).

Station 1 is located just below the bridge crossing the National Turnpike. The stream bottom is comprised of bricks, concrete chunks and other
types of solid rubble that is often times prevalent in streams that are
associated with human population centers. We sampled the bottom by holding
the dip net downstream, as we overtime some of the in-stream objects.

Also, we probed the vegetation overhanging in the stream to dislodge any
epifauna present.

Station 2, which we considered as a control station, was located on an unnamed tributary to Wilson Creek, which drains the eaterly slope of the South Park Hills, which are part of the County park systems. We sampled a small segment of the stream just upstream from the bridge on South Park Road and Hornback Road.

Station 3 was located on Wilson Creek approximately three miles downstream from the Valley. The creek had eroded a bed 6-8 feet below the level of the flood plain. The bed was composed of construction material rubble and alluvial clay deposits that was riddled with crayfish burrows.

Station 4 was located downstream from the Valley at the boundry of the county forest. Just upstream from this station, was a small dam that had been installed as part of the instream treatment system. Water was being pumped from behind the dam into a fountain manifold that had been installed

upland. The stream water was aerated to release volatile organics then allowed to trickle over the ground surface back to Wilson Creek.

Station 5 was located on a small stream that originated in the golf course located upland from the Valley. The stream was approximately two feet wide and .25 feet deep at this station. The flow as low but constant.

Creek approximately .25 miles downstream.

This feeder stream was approximately one foot wide and .5 feet deep. The flow was slight and formed ripples and pools as the stream flowed down the hillside. The stream bottom was composed of sand and gravel, with occasional clumps of allochonous materials, it dead leaves and twigs. This station was considered as another control as it drained a water shed that was within the county forest with little

## STATE OF KENTUCKY SAMPLING

On March 6th, a meeting was held at the Command Post attended by the following:

#### NAME

Robert Ware
Robert Logan
Robert Bay
Waynon Johnson
Don Schultz
John Gilbert
Royal Nadeau

#### AFFILIATION

Kentucky Div. of Water Quality Kentucky Div. of Water Quality U.S. Fish and Wildlife Service U.S. Fish and Wildlife Service USEPA— Region IV USEPA— ERT USEPA— ERT

The primary purpose of this meeting was to recommend a sampling survey that would generate the types of environmental information for the OSC, Jack Stonebreaker.

The consensus of the group was:

a. A sampling of Wilson Creek should be performed as soon as possible to determine the extent of impact relative to chemical and biological parameters.

- b. That some acute toxicity tests be performed on water collected from specific locations on Wilson Creek and from the "collection pond."

  The emphasis was not to determine a TL50 so much as to note any absolute toxic effects on the test organisms.
- c. That chemical analysis be performed on tissues of aquatic organisms collected from Wilson Creek to determine possible biomass contamination of specific organisms compounds originating on the Valley.

At this meeting the State of Kentucky representatives indicated that they would collect water and biological samples which would then be shipped to Region IV's Athens Laboratory for analyses.

On March 8th, a sampling team constituting of EPA-ERT and State of Kentucky personnel took water and sediment core samples from the same locations as identified for the biological evaluation sampling effort. In addition, water and sediments were collected from an abandoned settling pond within the Valley. Water for bioassays were collected from Station 02, 04 and 08 (collection pit). These samples were then transported to Athens under ice to the Athens Laboratory by Region IV personnel.

On March)9th, benthic organisms (mostly crayfish) and fish (minnows) were collected by State of Kentucky personnel near the same sampling location as previously identified (see Figure 3). These organisms were shipped to Athens under ice for chemical analysis of their tissues.

On April 16-19, 1979, State of Kentucky personnel collected additional biological specimens for chemical analysis. The samples consisted mainly of fish collected from Wilson Creek and some additional downstream station in Bee Lick Creek, Northern and Southern Ditch, and Pond Creek. The fish were collected via electroshocking and hoop nets, then sent on ice to Athens for chemical analysis.

# ANALYTICAL METHODOLOGY

Upon receipt by EPA Region IV Surveillance and Analysis Division's Laboratory, the samples were processed and analyzed according to acceptable analytical procedures.

# WATER

Water samples were analyzed by a modified procedure which has been devised for Priority Pollutants (revised April 1977).

# SEDIMENT

The sediment samples were first processed in a high speed mechanical deward and interest to break the large soil chunks to a uniform sediment particle size. These here the large were extracted with acetone/hexane at a 1:1 mixture, then analyzed in a Gas Chromatograph/ Mass Spectrograph.

#### TISSUE

Tissue analyses were performed at the University of Georgia under contract to EPA-Region IV.

Sample cleanup was accomplished through an automated cleanup device which utilized gel permeation. This procedure separates and sorts according to molecule size. Extraction was according to the FDA Pesticide Analytical Method as modified to accompdate the gel permeation cleanup technique.

# RESULTS AND DISCUSSION

The extensive environmental sampling and analytical effort revealed the presence of a vast array of inorganic and organic substances in the Valley proper and the adjacent Wilson Creek.

The substances of main concern for this report are the organics, as it is these compounds that information is most scarce. Therefore, more attention will be directed to these substances than the inorganic constituents,

ra

mos :

es t

-

# A.L. TAYLOR SITE

Station AT-7 was a settling impoundment which had been constructed by win.

Mr. Taylor to hold wastes emptied from durms that were set turned on the the upland slope of the pond. These analyses revealed a large number of solvents, ketones, pthalates and other organic compounds which are used or associated with paint and plastic industries.

The results of the sediment analyses revealed fewer number of organic compounds than the water samples collected from the same area within the Valley. Station AT-4 sediment samples revealed the largest number of compounds (28 residuation distinct identifiable compounds). Aroclor 1254 and 1260 were among the compounds identified from this station along with several solvent type wire compounds.

#### WILSON CREEK WATERSHED

The topographical location of the Valley is at the top of the watershed is a of Wilson Creek. This position also affects the hydrological gradient. The the informal stream flow in Wilson Creek adjacent to the Valley is low and subject the

to fruction from seasonal storm and melt water contribution. The stream bottom in these upper reaches mainly consists of sand and some clay deposits. These materials are the results of erosion that has carried the materials into the stream from adjacent upland area.

Wilson Creek does not have a USGS gaging station, therefore, there are no historical stream flow data. There is flow in the stream bed throughout the year, although very slight during dry periods.

#### WATER

The water samples collected on March 8th from Wilson Creek only revealed nine identifiable organic compounds with bis (2-ethyl-hexyl) phthalate occurring at Station 02, 03 04 with station 04 having the largest concentration.

An independent laboratory (Environmental Consultants) analyzed a water sample collected slightly downstream from the Valley. It contained methyl pentanone, ethyl benzene and xylene in trace amounts.

In another water sample analyzed by the same laboratory on the same day (March 4), xylene compounds were found in trace amounts.

On a separate sampling run performed by the State of Kentucky in (see 7 is use 5)
early February A, water samples collected just downstream from the Valley contained a vast array of organic compounds (38 identifiable compounds. See Table 2). Many of these were present at all of the station sampled in Wilson Creek. Most of the compounds occurring in the water on this sampling date are ones that are associated with paints and solvents. The water samples from the station closest to the Valley, contained elevated levels of methyl ethyl ketone and methyl isobutyl ketone.

#### SEDIMENTS

The sediment samples collected by the State of Kentucky in Wilson Creek were contaminated with trace amounts of organic compounds (19 identified compounds). Aroclor 1254 and 1260 were present at all station in Wilson Creek in above trace amounts.

# ECOLOGICAL EVALUATIONS

The results of the biological surveys of Wilson Creek proper performed by EPA biologists and State of Kentucky biologists are outlined in Table 3.

Only those station that were sampled by both groups are listed. The observations made by each are similiar and tend to collaborate each others results.

Some of the stations sampled by each group were not synonimous. For example, Station 04 through 06 of the Abiological survey were closer to the Valley while Station 04-08 of the State survey were donwstream outside of Wilson Creek groper. (TABLE 4).

The observations of the EPA survey for these remaining stations are listed in Table .

The populations of aquatic invertebrates living in Wilson Creek belong to those groups of organisms that are usually found in stream through the physical/chemical features. Many of these organism types can occur in streams that have been disturbed by man's activities. Therefore, their presence in streams is not necessarily indicative of stress being applied to aquatic ecosystem by the contaminants emanating from the Valley.

The presence of the macroinvertebrate species observed indicate that toxic compounds occurring in the water and sediments of Wilson Creek are not having a catastrophic deleterious impact on these populations.

More likely, the impact to the biological resource living in Wilson Creek is in the form of sublethal, low level contamination of biomass. This biomass

contamination is present within a variety of species forming the food web as indicated by the presence of certain organic contaminants, i.e., Aroclor 1254 and 1260 in the fish and macroinvertebrate tissues. It turns out that PCB's may not be the most appropriate organic compounds to show the impact of the Valley on these biological resources as there is not clear pattern to the tissue levels of PCB's that correlates to the location of the Valley relative to the sampling locations. Out of the 126 identifiable organic compounds found in the Valley, many have a potential for bioaccumulation as indicated from the Log P values (Table 5).

Log P is an expression of octanol/water partition coefficients which essentially indicate the solubility of a substance in a cample relative to its solubility in water. It is expressed as a ratio, for example, a Log P, a value of 4.0 indicates that the material is 10,000 times more soluble in octanol than water.

A substance that is soluble in octanol is considered to be potentially soluble in some of the biochemicals and that present in living systems. Once incorporated into living systems, these same materials can interfer with normal physiological functions, accumulate in certain tissues and be passed on through the food web to the next trophic level species.

Published biological concentration factors (BCF) that have been determined for specific organic chemicals, other than halogenated pesticides, are scarce indeed. As a result, the impact of many of these compounds in unknown or not well understood. This is the case for many of the compounds present in the Valley.

The presence of the abundant population of amphipods and ispods are indicative of allochthonous based food webs, being that these organisms utilize dead leaves and similar organic materials for food. These organisms are then preyed upon by more moible species as a food source, i.e., fish present. It is not

surprising to find this type of energy flow or food web present in a creek like Wilson Creek which flows through the heavily wooded area near the Valley.

An interesting aspect regarding the fate of contaminants are large molecule organics, with various degrees of water solubility. Those that are rather insoluble are sorbed onto the particulate or suspended matter, i.e., eroded soil particles, dead leaves and bark pieces, etc. Some of these contaminants are deposited in the stream bottoms or in shallow shoal areas.

Some are incorporated into the food webs when consumed by the indigenous populations feeding upon the allochthonous materials or bottom deposits.

It so happens that the organic scan performed on the tissues collected were for PCB's and pesticide compounds water analyses did not indicate the presence of halogenated pesticides in the Valley, therefore, it is not surprising that these type compounds were not detected in the organisms analyzed. These same analyses do indicate that there are other compounds, some of which are known toxics, i.e., Priority Pollutants, etc., that are present in the Valley. These are the types of compounds that if incorporated into the biological resouts could seriously affect the utility of these resources. This phenomena has been well documented for Mirex in Lake Ontario and Kepone in the James River. In both cases, the biological resouts affected were determined to be unfit for consumption or any other type of exploitation.

Since the tissue extracts were only analyzed for PCB's and halogenated pesticides, other organic compounds may be present in the tissues, therefore, it may be wise to expose these extracts to GC/MS examination. GC/MS would reveal the presence of other compounds, i.e., PNA's that might be present.

#### MODE OF CONTAMINATION

The main mechanism of environmental contamination to the aquatic resources is via surface and just below the surface runoff into the Wilson Creek watershed. This was obvious in the initial EPA investigation and actually triggered the activation of a 311(k) cleanup by Region IV.

Surface water containing oils were sighted at this time, however, the underlying, more serious concern related to the numerous potentially toxic compounds that were in the drums.

easily quantified. Surface runoff is the runoff is upon rainfall and snowfall.

An examination of the Climatological Data Record for the Louisville Weather

Office of the National Weather Service located at Standiford Field Airport revealed for the spring in amounts and types, for the spring when most of the sampling and mitigation activities were occurring.

The actual area of the Valley containing drums was about 10.1 acres as determined from the USGS Topographic Map (Brooks, Kentucky Quadrangle). The in the Valley are relatively impervious, therefore, most of the precipitation falling on these 10 acres will end up as runoff.

The collection pond and collection ditches were designed to entrap most of the runoff emanating from the drums storage area. According to the latest estimates, the effluent flow from the in-place treatment system is approximately 227 liters/min (60 gals/min). The system is operated for ten days out of a month during the summer, thus, the discharge from the Valley for this summer was approximately 327,059.5 liters/day. Negating evaporation from the collection pond, the discharge volume would approximate the amount of water being contributed to Wilson Creek from the Valley, prior to the mitigation procedures.

Knowing the discharge rate and the concentrations of materials in this discharge, it is obvious that significant amounts of toxics were being con-

tributed to Wilson Creek from the Valley. For example, Diethyl Phthalate, which is one of the Priority Pollutants, would be discharged from the Valley at the rate of 8.5 grams/day for the summer months; perhaps more for the months when flow would be greater.



# MAN'S USE OF CHEMICALS FOUND IN THE VALLEY

It was difficult, actually impossible, to determine the origin of all the compounds found in the Valley being that records of the origin of the drums were not available. Therefore, we resorted to performing a literature search on each identified compound in hopes of finding its use within our present society. Table & represents the results of this effort. The categories listed are the more common uses of organic compounds found in the Valley. As expected, not all the compounds identified could be catagloged, these are indicated by the blank space. It is a majority of the 126 compounds found are used as solvents or associated with plastics or paint manufacturing.

The rest are distributed rather evenly throughout the other uses listed.

TOXIC EFFECTS OR ORGANIC EFFECTS FOUND IN THE VALLEY.


When the results of the initial environmental sampling revealed the extent and types of organic compounds in the Valley, a literature search was performed to determine the known effects of these compounds. Table 7 categories the effects according to the effects documented in the Registry of Toxic Effects (Fairchild, et al, 1977).


Out of the 142 compounds, 40 are included in the Human Effects section of the Registry of Toxic Effects, 27 are in the Carcinogenic Effects Section, and six are in the Teratogen Effects Section of the Registry.


In order for a substance to be life the Carcinogenic Effects Section, there must be documented evidence that Malignant tissues are produced by the substance in the body of the test animal. For a material to be included on the Teratogenic Effects, it must be documented that changes in offspring are produced but are not transmissione to their offspring. Those substances that are listed in the Human Effects have been found to have produced responses that is considered deleterious in some manner by the panel of experts that review and screen these materials.

#### Literature Cited and Data Sources

- 1. McLeese, D.W.; V. Zitko and M.R. Peterson 1979. Structure-Lethality relationships for phenols, anilines and other aromatic compounds in shrimp and clams. CHEMOSPERE 2: pp 53-57
- 2. Veith, Gilman D. and R.T. Morris 1978. A rapid Method of estimating Log P for organic chemicals. EPA-600/3-78-049. 16 pgs.
- 3. Standen, Anthony (editor) 1964. Encyclopedia of Chemical Technology. Vols. 1-22, Second Edition. John Wiley and Sons, Inc.
- 4. Stecher, Paul G. (editor) 1968. The Merck Index, Eighth Edition Merck and Company, Inc.
- Fairchild, Edward J and R.J. Lewis and R.L. Tatken (editors) 1977.
   Registry of Toxic Effects of Chemical Substances, Vol. 2.
   U.S. Department of Health, Education and Welfare, DHEW, Publ. #78-104-A









#### OWLEDGEMENT

This report would not be possible without the cooperative efforts of crous individuals associated with Region IV's Emergency Response Branch, ar Surveillance Branch, Technical Support Branch and Ecology Branch plus Environmental Response Team and the Kentucky Division of Water Resources.

1	-						•		1.
(לפפ) פרוסקיים	MATER SER		SEPIMBNIT WATER	7-3		SEDIMO	AT	-5 AT	1 AT-
Servoic Acid 1'		22.22	36/14/6/	JESTAER	TONIER.	10000	1-25	1-10	10-50
" methyl dentale Add "1 (somets)			/0-/06	==	10-100		1-25	1-10	10-50
Co Alayl Benzoic Acid		7.9	10-100	==	1-10			1-10	1-10
tinidentified compounds /-/0 ppb	,	7	18	-	16	37	1-25	20	/-10
Wintentified compounds 10 - 100 pph (Viene (2 isomers)			260		120	330	470	62	300
Mechyl Pencandi Mechyl Cyclopencandi				==					7.7
Sucoxyechanol				==	<b></b>				77
Techni hydroxy Pentanone  L- Alkvi Benzene () (somets)			24 64		170	10-100	7.2	2.8 5.7	200 27
Disucovvectanol (2 (vomets)			65	==	150		86	15	60
Mechyl Styrene C. Akayl denzene (5 isomers)			22		111	10-100	26		28
a,a Dimethyl Sendere Pethanol Ch Alayl Styrene			5.6	-				19	7.3
Isophorane 2/				-				15	7.3
Thenoxy Methyl Unitare					1	1-10	-	144	15
Chiprophenoxypropingi Dimethyl Phtmatate -/				<b>—</b> —	8.7		9.6	1 /3	
Prenoxyptopaneu12: Tetraoxagodecane		220	4.6		20		7.2	2.7	57
Methyl Propyl Ester of Benzoic	9.4	17	7.5		4.9		29	26	3.7
Acid Pentaoxapentadecane		320	27	-	4.9		36	7.7	3.7
Bis (2-echylhexyl) Phinalace 2/	13	45			1	120			1070
Phenol 2/				+==	741	Tel	R/	THI	120
Tetnyl frenci Co Alkyl Phenol (3 isorers)			1-10	==	1-10	7.3	1-25	1-10	10-50
CA ALAVI FRENOL			96	==	-		1-25	10-180	1-12
Directive Promano.				<del> </del>	10 400	0	1-25	3F0 (#0	10-50
france-t, 2-dichterstrytene 2/	745 PK	1025 745	75	FK.025		8	36 .	745	6.4
CCAVI Senzene ./			340	-29			180		31
Cricolorocriflueroechane				6.0			Vas		5-7
Prechyl Ethyl Ketone -/			730	==			¥10		27
I-hexanol				==	78		1.6	TUT	1530
l-methyl-1-promanol Tetrachioroethane	6.4	12	12	<u> </u>	745 5.7		12	62	5.2
I-butanol I-bexanol			3.73 8 7 7				T45		765
i.l-dichloroetnine			3H 8 6-01 F	<b></b>	><5		745		745
Vinvl chloride			200		Tes				745
Methylene Chiorice 1/ Trimethyl Cyclonexanone	6.6	7.9	5-3				1.4	7.4	
Methyl Benzene Methanol Carponic Acid, Busyl Phenyl Ester				+==				3.5	
Propoxy Butane Di-n-putyi Phinalace 1/	39	7.5	3P 5:6	1-10	10		9-6	1.2	
Fluoranthene 2/ Dimetryl Phenol (3 isomers)					¥.9		7.2	747	
Hexanol Acid			1-10				1-25	1-10	
Pentachiorophenol 1/-1/ Niphthalene 2/			1.9	1//:	74/		4.5	. 4.3	
Campnor 2/				-			9,4		
Ch Alkyl Benzolt Atia metnyl esters of Ethyl mexamolt Acid				1-10	1-10		1-25		
Benzene Propanic Acid [Heptanol.							1725		
neotanone Incene				-			220		
Methyl (methy ethanol) cyclohexanol							42		
Terpineneol Tsophorone 2/			: 179	==	20		113	- ;   -	
Cutnoline Mednyl Daphinalene			- 77			4-11	9-6		
Acenaonchene =/						1-10	9.6		
Dipensoturan Beta-naphtnomitrile							W.F		
Phenanthrene, Actional L						/	7.2		
isutvi l'otnyi Benzere sulfonamide				==		1-10	0.7		
Pyrene !/				_			7.2		
Sicory Propanol Ca Alkyl Styrene			4.7		6.2				
Pachyl (Proposylovy) Ethary Probinol			24		12				
Totra metnyi Pentanone					11				
Diputyl Einer			5:6		5.2				
ic Alayi Benzene			1,4	<del> </del>		1-10			
'etnyl mestensne		45	2.5			1-10			
Icinechyl Pencandiol Icinechyl nexune	6.7 4.8								
recove Pronoxy Pronunce	5.7 3.2			1					
Stoay Fronoxy Methane									
Stretryl Ethyl Benzolc Acid									
Tethone Tethoxymethylethoxy Propanol					-				
Stausvioky rendanda			-:-					·	
**ITYL Pronancic Acid									
- Cannot Asia									
Tit Honors				==		1-1-			
						1-10			
						1-10			
forthernyl phenylcinyarpingene						1-10			
NATUR Prenunchasere						1-10			
147712 AC.1						7.4			
ACCOUNT ACCO				==		3.2			
encade and a control of a contr						70			
- residenting and a control of the c						370 200	-		
		1 1 1	1. 3. d f		in the state of				TARREST STATE
			and the same of th	A CHEN LA	. Carried and			- to the second of the second	· British Street

		· .	•						The second second
				-002	We	-003	WC	-004	Table 2.
10×20/00 (222)	WATER	ונש ביינש?	WATER	CEPVALEA	WATER	Fame	VT WATER		
Benroic Acid I/	14	ļ	1-10		1-15	<del> </del>		725	
In Alavi Benzolc Acid			1-10			$\pm =$			
Cettor Senzene sufferante	P. 7_		1-10		1-15	$\pm =$			
Unidentified compounds /-/0	13		12		15	<u> </u>	18	<u> </u>	
(Viene (2 isoners) (Petryl Pentanol	43		68	-	22	F	1 /3		
Methyl Cyclopentanol Discrevi Pentanone						$\bot =$			
Butakvethinal	- V-					<b>+</b> ==			
Pechyl nydroxy dentanone - Alkyl Benzene (3 isomers)	9.5°	<u> </u>	9			上	1.0	1	
Diputatione (2 (sames)	5.7	<u> </u>	7	7<3	3.3	==	1.0	7 35	
Tetavi Styrene C; Aimvi denzene (5 isomers)	9.6	<del> </del>	40		4.2	F==	1.8	<del></del>	
1.a Dimethyl Senzene Pethanol To Appl Styrene	A 3	-	18	-	F.V	<b>T</b>	6.3		
tatak Sebican Socaras,					<u> </u>	‡ <u>==</u>	-		
Isochorone 2/ En-nosy Meshyl Detrane				<del> </del>		<u> </u>			
Interconenceveropanul				<del> </del>		<u> </u>		<u> </u>	
Othernyl Phinaiate _/	2.2				<del> </del>	$\vdash$	- <del>ļ</del> -		
Tecraoxadodecane Diechyl Phonalage 2/	1.2	<u> </u>	16		2,2	<del>                                     </del>	1 1.8		
Methyl Promyl Ester of Benzoic Ach							ii .		
Pencaoxapentadecane	9,3		30	<u> </u>	6.2	==	5.1		
Bis (2-echythexyt) Phchaiace 2/ Pechyt ducanoic Acid	741	745	1-10	·		<del> </del>	- <u>1</u>		
Phenot 2/	74/	745	1-10		741	⊢	5<1	+	
[ethy] rhenoi [r Aikvi Phenoi (3 isomers)		Y < 5	1-10			$\vdash$		725	
Di olevi daenoi Dinyi mekundic heid	93			745	2.5	二二	ļi .		
Ospecant Propanol Phonatic Acid	2,3		<b>—</b>	<del>                                     </del>	1-15	<b>=</b> =	2.2	1	
teams-1, I-dicalarat-vivae 2/	43		114	<del>                                     </del>	į	<u> </u>	r H	<u> </u>	
Columne L/ L/ Start Sensene L/		<u> </u>	3/	<del> </del>	<del>}</del>		-	<del> </del>	
Cotone Crichlorotrifluoroethane	77.56		9,000		3	$\vdash =$	1	-	
rechyt Ethyt Kecome ur Sexide	7500		6300			$\vdash =$	-		
Totavi Isobutvi Ketone 2/	2600		12,000			$\vdash =$	<u> </u>		
l-metavi-l-propandi							1		
Cecrachioroethane C-bucanai			24		ÿ 2		<u> </u>		
l-hexanol L.l-dishlorogenàne	745		3/4	777. ***	1 0 1-2-			F	
1.1-dichloroechina			-	7 86	64 5		1		Ŭ .
Dimetavi disultice Metaviene Chiptice 2/	58		7 63				<u> </u>	<del> </del>	
Trimethyi Cyclonexanone	38	<del></del>	52_			<u> </u>			
Mechyl Beniene Néchanol Carponic Acid, Butvi Enemyl Escer					<u> </u>	上—	r fi		<u> </u>
Propoxy Sucane Di-n-pucyl Probalice 1/				ļ <u>.</u>	<u> </u>	<u> </u>	1	<u> </u>	
Fluoranchene 2/ Dimethyl Phenol (3 isomers)	721			1					
Hexanom Acid Evaroxy Mechaxy Benzene		745					-	74.5 74.5	
Pentachiorognenol 1/ 1/			3.6	<u> </u>	1				
i.ifrichiosoethine			216		7610		721.0		
Cabbnor 2/ Ca Alkyl Bensold Acid				<del> </del>					
Pecnyl esters of Sthyl dexample Acid Senzeme Propanic Acid	سے ق	725	100		16	<u> </u>	/ / 3	745	:
Feotanol Feotanone	62-		108	<del>  </del>	1 15	<u> </u>	35	<del>                                     </del>	
incere Rechyl (methy echanol) cyclohexanol							1		
i-Terpineol Terpinene-4-ol					<u> </u>				
isophorone 2/									
Muinoline Mechya Bapachalene							<u> </u>		
Atendontheme 4/ Stophosfuran							l l		
Baca-maintnomitrile Fluorene						<u> </u>	<u> </u>		
Tributol Escer of Prosphoric Acid Phenanthrene/Anthracone 44						<b>—</b> —	8		
Sucry Jethyl Sendene Susionamide						F==	b .		
Gethy, Octanone	1.3					==	<del> </del>	735	
Sitory Proganol Sa Alayi Styrone			5.0		74.0		1		
TITANKA ELUOKA ELUTUA!	7.2		6/		22		17.0		
vicatine Staecayi caenei							<u> </u>		
Tours methyl Pentanone >- Tiputyl Ether									
ls Aikvi Benzene		#							
.) mist:Vrene			<del></del>				<u> </u>		
Corecavi densandioi		- H							
Inthetoxi nexune Tetoxi Pronoky Pronanol									
- GEORY Espoury reordings SEGRY Espoury Mespour							ļ		
retrie cores de Adesv.	29						ļ		
Tresnot Echyl Bensold Asid	7.2	7 < 5							
rethykymechylethoky Probanal	4.9								
t nyanathaenune	1.9	N N	5-		2.7		213		
retryl Problemote Attid		B	1-10		1-15		41		
-:3-2:1 \:::5 -:20023001			1-10				ļ		
♥/. (icerol									
ATTOAMACTIAL TROLLEDS									
- Austroomenoù - Aust									
cometric medvicovaratraene cassar luie and lued 2		./3		5.0		.0097		. 52	
Nivol rhemanintene Nivolenza inimalace i/				7.0		====	ļ		
.1.** -11 -2114									
		———F							
Frombia Walt									
							ļI		
77:7011 1012 #707012 4111 [-#710015 #112]		755		745				7 4 3	

# TABLE 3. MACPOINVERTEBRATES FROM WILSON CREEK WATERSHED

State of Kentucky (March 19)

Station 01. (2.8 miles below Valley)

Amphipoda

Gammarus sp.

Isopoda

Lirceus sp.

Decopoda

Cambarus sp.

Station 02. (Fairdale Raod Bridge Area)

Amphipoda

Gammarus sp.

Isopoda

Lirceus sp.

Decopoda

Cambarus sp.

Mollusca

Gastropoda

Physa sp.

Trichoptera

Limnephilidae

EPA (March 7)

Station Ol. (National Turnpike Bridge)

Amphipoda

Gammarus sp.

Isopoda

probably Lirceus sp.

Tipulidae \_

Tipula sp.

Station 02. (Control Unnamed Tributary to Wilson Creek)

Amphipoda

probably Gammarus sp.

Isopoda

probably Lirceus sp.

Decopoda

probably Cambarus sp. (juveniles)

: Platyhelminthes

Turbellarid flatworms

Mollusca

Gastropoda -

Probably Physa sp. (snails)

Ephemeroptera

Unidentified

Trichoptera

Unidentified

# EPA MACROINVERTEBRATE SURVEY March 7, 1979

#### STATION

04
Just downstream from aerator

05
Just downstream from Golf Course
pond outlet upstream from Valley



06
Riffle-pool area on feeder stream into Wilson Creek

#### OBSERVATIONS

Isopoda probably Lirceus sp.

Amphipoda probably Gammarus sp.

Oligochaeta unidentified

Isopoda probably Lirceus sp.

Decopoda probably Cambarus sp.

Amphibia salamander egg masses

Amphipoda probably Gammarus sp.

Isopoda probably Lirceus sp.

		•		TAB	LE S.
(בפר) באינים	46 P	BCF			2000年1月1日 - 1000年1月1日 - 1000年1日 - 1000年
Sensote Acid 1/			<u> </u>		
C. Alevi Benanic Acia C. Alevi Benanic Acia (actu) Benanic Alia					
1 Unicentified compounds					
(Viene (2 isomers)	/,25				
Methyl Cyclosentanol Dimethyl Pentangne Butosyethanol					
Techni nudrosu fentanone					
Othucosymechanol (2 (somets)	97, 95				
A.a Directvi Benzene Mechanul					
Isopharone 1/					
Cheny Siner					
Chiorognemosypropanol Cimetnyl Phinaiate 2/ Champaygropanedici	2.//				
Tecraaxadadecane Diernyl Phobalage 3/ Mechyl Propyl Ester of Benzolc	7.15				
Penragapentageane					
31s (2-ethylhesyl) Phthalate 2/					
Phenol 2/ Phenol 2/ Technic chrosi	1.49 1.64	·			<u> </u>
Co Algul Phenot () teamers)					
Directory Prominos					
Convers of all and a second sections of the second sections of the second secon					
School Control					
rethyl Sinyl Necome -/		5 - A	1 1 16		
"nenvi Isobucvi Ketone 1/  -herandi  -metavi-1-medanadi	45 26 98 44	2. T.			
letrachiornethane	364				
l'herangi i,i-dichiorpechane i,i-dichiorpechane	1.79				
Vinvi Chioride I/	1.77				
Methylene Chiorice 2/ Trimetnyl Cyclonexanone Methyl Senzene Methanol					
Proposy Butane					
Di-n-pucyl Phohalace 1/ Fluoranchene 2/ Dimecnyl Phenol (3 isomers)	2.36		<u> </u>		
Hexano Acia Hysroxy Methoxy Senzene	1.94, 1.90				
Pentachidroonenol 1/ 1/ Naonthalene 4/ Li,-trichidroethane	3.76, 3.72	7,70 1.99 4.26			
Campnor 1/					
Mechyl esters of Ethyl Mexamold Acid Benzene Fromanic Acid Heptamol					
reptanoné incene	2,4,2_				
Methyl (sethy ethanol) cyclohexanol a-Terpineol Terpineool					
isaphacane 4/					
Metnyl (danthalens Adenabninens s/ Olionzafuran	3.93_				
Beta-magninomitrile					
Tributel Ester of Phosonosic Acid Phenanthrene/Anthronens 44 Sutvi Pothyl bensene autonamide					
Pyrone 4/ Methyl Uctanone					
Sitory Probanci Co Alkyl Styrene Setayl (Francisco) Stadey Probanci					
Survory Stadky Espanol					
Totta metnyi Pentanone Jibutyi Einer					
To Alkyl denzene					
it nine, ilvedia letnyi medcendna letnyi medcendna					
Finetovi Perane Petovi Pronew Promot					
TISCHAN PENDANA PENDANA TISCHAN PENDANA PENDANA					
incoming loss incoming femal Bensold Acid	1.40				
Tothore Tothokimethylethoxy Prepanal Loutyloxy Prepanal					
i nysrusnaenune etnyk Pennamoko Askd					
corpt pactamote Acta					
-rrocinci Tilinorpi Tilinopi					<u></u>
restrongnoses					
Control Publican  Control Inerwisinvaroindene  Control Duke and Loui Lo					
12-71 Penanthrene					
141911 A212 11011 1111 11011 1111					
TITA-companie acts Tita-companie acts Literarchi acts					
+ BCF = Die CONCENTERT	- FACTOR	en en jaron eta	44 M (200	HATEL TO SERVE AND A SERVE AND	
= Loc P = Voices from Es				BOOK TO THE SECOND SECO	· The Control of the

Column   C	e e e e e e e e e e e e e e e e e e e			. •,	٠.							TA	BLE	6.
Company   Comp	(לכבר) באומצים	PAINT	SOLVENT	PLASTICS	P'SERATA	SE MATINE	LE PARE	PERFUSA	DYE	1411 CEPTIC	१ र र र र र	אבן ופסרונות	-	
Section of the control of the contro	ethy: Seniold Adid (C isomets)						<u></u>							
	CTAAL SAUSOFC, VCTT						<u> </u>	-			<u> </u>		<del>                                     </del>	+=
Comment   Comm	Unicentified compounds													
EATER CHARGES AND	ernet Pentanol		+-				<del></del> =_							
CONTRACTOR MATERIAL STATES AND	) Leetnyl Pentanone						==					<del> </del>	<del> </del>	<del>                                     </del>
	echyl hydraxy Pencanone										<u> </u>	-		+
STONE CHANGES AND STONE CHANGE	- 2 - 2 7 e 4 a none		<b>+</b>											
Treatment of the control of the cont	ecnyl Styrene Alay denzene (5 Liozera)			+										
AND THE CONTROL OF TH	Livil Styrene		+		<u> </u>							<u> </u>		<del> </del>
STATE OF THE STATE	soonorane 2/													-
TOTAL DESIGNATION OF THE PROPERTY OF THE PROPE	nonvi Ether			t										
ACCOUNTS AND	zenexyptopaneu.cl		+								-		-	+
Edition Section 2	terryt Chematage 2/													<del>                                     </del>
A CONTROLLED AND ADMINISTRATION OF THE PROPERTY OF THE PROPERY OF THE PROPERTY OF THE PROPERTY OF THE PROPERTY OF THE PROPERTY	Acid encapsapentadecane			+			<u> </u>							+-
AND A STATE PROBEST OF THE PROPERTY OF THE PRO	1s (2-ethylhexyl) Phthalace 2/ ethyl sutanoic Acid					+								
A LANCE PROBLEM CONTROL OF THE PROBLEM CONTRO	henoi 2/		+					+				I		
STATE CONTROLLED AND ADMINISTRATE AND AD	Alkvi Phenoi (3 isomers)													-
SAME ALL MANUAL PROPERTY OF THE PROPERTY OF TH	my: metanoic Acid												<u> </u>	<del>                                     </del>
TOTAL STATE OF THE PROPERTY OF	nthalle Acid													
TOTAL CONTINUES AND THE PROPERTY OF THE PROPER	invi Sentene i/		+	+				सामित्र प्र	17359					$\pm$
TOTAL SCORE 3. P. T.	richlorotrifluoroethane		+					الانطا				· ·		
SCHOOL STATES AND STAT	tane		+				==		<del>_</del>					=
SMISSION OF THE PROPERTY OF TH	necovi-l-scommol									· · +····				
Comment   Comm	-butanol							<del>_</del>						
price control of the	1-dichiorpechane													<del> </del>
AND STATE OF THE PROPERTY OF T	invi chiorida I/.													<del>                                     </del>
ASSOCIATION AND ASSOCIATION AS	chylene Chloride 2/		<del></del>											
	itponic Acid, Butvi Phenvi Ester													
INCOME PROPRIETA  PROP	i-n-putvi Phinalate 1/					•								<del> </del>
Section Microse Section  SECTION FOR CONTRACT  SECTION FOR CONTRAC	imecnyl Phenoi () isomers)									+	<u>+</u>			
	encachiorophenol 1/ 1/				4-									
ALEXT DEFORM ASSAULT	trichiocoethane	<del>=</del> =			1				7					<del> </del>
TOTAL PROPRIES AND STATES OF THE PROPRIES OF T	Alkvi Benzoic Acid convi escers of Echvi Hexanoic Acid		<b>\</b>						_+_		+			
SCIENT	enzene Propanic Acid		+											
	cene							+						
	Terpineol	==			<del></del>		<u> </u>	_+_					-	
Control   Cont	concrone 2/													<u> </u>
Command   Comm	covi Rapochalene		_ †	+										
	ca-magacmonitrile						$=\exists$							
	routyl Eater of Phosphoric Acid												<del> </del>	-
Control   Cont	tvi Nethvi penzena bultonamida						==						<del>] -</del>	
test in fromework female  Jamiline	Cory Propanol	==												
Intentive (CORDI)	thyi (Tropoxyloxy) Linoxy Prooundl						===						+	
Nierl Benerer	Setavi (Senoi												-	
A	cra methyl Pentanone	===	+				$= \exists$							
	WYKAT LUGUOT						$= \exists$		I			*	+	
Comment   Comm	thy: destensing		+				==						-	
ACCOUNT PRODUCT   ACCOUNT   ACCOUN	inecovi mexine covi Persoxy Personal							-1:						
Intervit Ethyl Benrott Acid	GEORY Proposy Pechane													
# Mone # # # # # # # # # # # # # # # # # # #	95 19 (15 <u>15 1</u>						===	-						<u> </u>
1	rinong						= =							
MOVE REPLACED ACTS  **CONTINUENT	DAGIOTA LIGORAGE						$=$ $\pm$							
Process   Proc	thy: remtampic Adia						$= \exists$	+						
Interview of the control of the cont	12023071						$= \pm$	+		4-	<u> </u>			
SIGNATURE STATEMENT STATEM	:nen:>0:uF1000e													
Intervit   Intervitiva   Int	-seraph-dras/ -culture abbetaran						$\equiv$							
### SECTION AND AND AND AND AND AND AND AND AND AN	Stiur (11)+ And (12) (7						=							
Profit   P	Cri tensi inthalace ir						$= \exists$							
Principle Alia   Ministry page Alia   Ministry pa	737516 3553						==							
	Immangio Asia Entremandio asia						==							
Therefore and the second secon	7:1-cer:7316 A:15						==	<del> </del>						
・ 「「「「」」」、「「」」、「」」、「」」、「」」、「」」、「」」、「」」、「」	132000000000000000000000000000000000000													

and the second section of the second section s